


10/36 Ware Street, FAIRFIELD
Phone: 9724 6864
Fax: 9724 0256

Dr. Diep Nguyen 204651 AX
MB.BS. FRACP, FAANMS
Specialist in Nuclear Medicine
Dr. Lyn Chan 209004AA
MB.BS. FRACP, FAANMS
Specialist in Nuclear Medicine

- | | | |
|--|--|---|
| <input type="checkbox"/> BONE DENSITY (DEXA) | <input type="checkbox"/> THYROID SPECT/CT | <input type="checkbox"/> Other Scan: |
| <input type="checkbox"/> BONE SPECT/CT | <input type="checkbox"/> PARA-THYROID SPECT/CT | <input type="checkbox"/> More Request Froms |
| <input type="checkbox"/> STRESS SESTAMIBI SPECT/CT | <input type="checkbox"/> Gallium SPECT/CT | |
| <input type="checkbox"/> GATED HEART POOL | <input type="checkbox"/> Hepatobilliary SPECT/CT | |
| <input type="checkbox"/> LUNG SCAN | <input type="checkbox"/> Renal DMSA/ Renal DTPA | |

BULK BILLING



Preparations - Please bring all previous scans

BMD (DEXA) Scan, Bone Scan, Gallium Scan, Lung Scan, Liver Scan, Gated Heart Pool Scan, Labelled Red Cell Liver Scan

- No preparation required

Thyroid Scan

- Ensure that you haven't had a CT with contrast injection in the last 2 weeks. You must also bring a thyroid ultrasound or blood test results to your appointment if any.

Stress Sestamibi Scan

- No caffeine for 24 hours before the test. This includes coffee, tea, chocolate, coca cola, herbal teas or decaffeinated drinks.
- Have a light breakfast on the morning of the test.
- If you're a diabetic, take half of your insulin or tablet on the morning of the test.
- Please bring all your heart and blood pressure medications with you as they can be taken once the exercise has concluded.
- Bring comfortable clothing.

Hepatobiliary Scan

- You must have something to eat 3 hours prior to the test and then fast until the test is completed.

Gastric Emptying Scan

- Do not have anything to eat or drink for 3 hours before the scan.

Labelled White Cell Scan

- If you are having the study for colitis, fast 3 hours before the test.

Gastro Intestinal Bleeding Scan

- You must fast 3 hours before the scan.

Colonic Transit Study

- Stay off any bowel medications including laxatives for 3 days before the test.

DTPA/DMSA Renal Scan

- Drink a minimum of 1 litre in the 1 hour before the scan.

APPOINTMENT DATE/TIME: